

AUTHOR INDEX

Bold type indicates authors of commentaries, editorials, and Resource Corners; (No.) indicates issue number.

- Ace Inhibitors in Diabetic Nephropathy
Trialist Group, 41
- Adams NP, 22
- Adkisson W, 51
- Akriviadis E, 4
- Alderman MH, 7
- Aldous MB, 31**
- Alejandro MM, 82
- Alpert JS, 113**
- Amin Z, 33
- Anti-Tumor Necrosis Factor Trial in
Rheumatoid Arthritis with Concomitant
Therapy Study Group, 3
- Apolinario PC, 71
- Appel LJ, 61
- Armstrong DG, 15
- Arnaud CD, 95
- Arroll B, 61**
- Arterial Revascularization Therapies Study
Group, 89
- Arvilommi H, 106
- Atallah AN, 11**
- Babikian V, 86
- Bailey-King S, 96
- Bartle B, 18**
- Baskerville N, 116
- Bates ER, 88**
- Bates SM, 110
- Bathon JM, 2
- Bauer TM, 75
- Bazelmans E, 47
- Beck C, 94
- Becker LA, 116**
- Beghe C, 76**
- Belknap S, 14**
- Benhamou Y, 53
- Bentall RP, 46
- Bentley E, 66
- Bernard GR, 81
- Bernsen RM, 24
- Bestall JB, 22
- Beta-Blocker Evaluation of Survival Trial
Investigators, 84
- Beyth RJ, 18
- Bhatt DL, 5
- Bilous RW, 15**
- Bjornsen SS, 45**
- Blacker B, 35
- Blanchard AR, 37**
- Bleijenberg G, 47
- Blood Pressure Lowering Treatment Trialists'
Collaboration, 6
- Bock SA, 35
- Borson S, 74
- Bossuyt PM, 109
- Botla R, 4
- Boulet LP, 44**
- Bridges-Webb C, 30**
- Briggs W, 4
- Brousseau L, 99
- Brown DL, 90
- Brun-Buisson C, 82**
- Büller HR, 16
- Burton M, 108**
- CALM Study Group, 9
- Campbell AJ, 100
- Campbell TF, 108
- Camus D, 27
- Canadian Task Force on Preventive Health
Care, 63
- Capsuto EG, 76
- Carewicz O, 43
- Carvedilol Prospective Randomized
Cumulative Survival Study Group, 85
- CAVATAS Investigators, 91
- Cello JP, 51
- Chan FK, 92
- Chang CJ, 90
- Chapman GD, 90**
- Chew DP, 5
- Chey WD, 53**
- Chiquette E, 11
- Chung S, 92
- CIDS Investigators, 113
- Clarfield AM, 94**
- Clark CR, 111
- Clarke PD, 27
- Coats AJ, 85
- Cohen H, 60
- Collaborative Group of the Primary
Prevention Project (PPP), 8
- Connolly SJ, 112, 113
- Conway DSG, 112**
- Cook DJ, 81**
- Cooksley GE, 52
- Cooper MG, 62
- Corley DA, 51
- Cowan FM, 64
- Craig JC, 62
- Croft AM, 68
- Crowther MA, 17
- Cullum NA, 55
- Cummings SR, 95**
- D'Ortho MP, 37
- Dans LF, 82
- Datta M, 67**
- Davenport J, 86
- de Jongste JC, 24
- Deeks J, 29
- Delaney BC, 114
- Dello JP, 51
- DeLoughery TG, 96**
- Denman AM, 2-3**
- Devlin N, 100
- Di Blasi Z, 98
- Diggle L, 29
- Donnan GA, 86**
- Doody RS, 94
- Douketis JD, 59**
- Duley L, 10
- Easter L, 61
- Ebrahim S, A-17 (No. 2), 60**
- Eckert K, 105
- Edwards RH, 46
- Eisenberg MJ, 34
- Ekeberg G, 12
- Elhendy A, 32
- Engelgau MM, 45
- Eriksson JG, 101
- Eriksson U, 83
- Ernst E, 98
- Espeland MA, 61
- Evans M, A-11-12 (No. 2)**
- Fann CS, 90
- Feagan BG, 25**
- Feder G, 66**
- Fehrenbach J, 75
- Feldman HM, 108
- Fennerty MB, 54
- Ferrer RL, 11
- Fielding K, 66
- Fife C, 76
- Finnish Diabetes Prevention Study Group, 101
- Fioretti PM, 32
- Fisher M, 9**
- Fjærtøft H, 12
- Fleischmann RM, 2
- Forster AJ, 117
- Forster J, 29**
- Fowler MB, 85
- French RS, 64
- Furukawa T, 102
- Gale T, 103**
- Gardner MM, 100
- Garner P, 27, 68**
- Garzon P, 34
- Gazmuri R, 117**
- Geleijnse ML, 32
- Gelenberg A, 103
- Georgiou A, 98
- Gerzmann JF, 77
- Gerzmann M, 77
- Ghahramani P, 88
- Gijssman H, 102**
- Gili L, 25
- Glasziou P, A-11 (No. 3)**
- Gluckman R, 8**
- Goeree R, 113
- Golish JA, 37**
- Gould MK, 73
- Grand'Maison A, 110
- Groeneveld PW, 5**
- Grupo de Estudio del SIDA 04/98, 26
- Gruson D, 56
- Gubareva LV, 42
- Haggard M, 108**
- Haidet P, 98**
- Haley CE, 42-43**
- Hannah ME, 10**

- Harkness E, 98
Hawkey CJ, 54
Hawley C, 103
 Hayden FG, 42
Haynes RB, A-12 (No. 3)
 Healy J, 33
 Heathcote EJ, 52
 Henderson-Smart D, 10
 Hernández-García C, 20
 Herxheimer A, 97
Hiatt WR, 109
 Hilbert G, 56
Hillegass WB, 90
 Hipolito RB, 71
 Hirsch AT, 5
Hirsch CH, 50, 74
Hlatky MA, 89
 Ho VA, 107
 Hochman JS, 57
Hodnett E, 105
Hoffman RM, 99
 Hogg W, 116
 Høgh B, 27
 Hohnloser SH, 14
Hotopf M, 28
 Howell CJ, 65
Hudson MP, 57
 Humphries KH, 112
 Hurskainen R, 36
 Hutten BA, 16
- Indredavik B, 12
 Ioannidis JP, 115
 Israel EG, 72
- Jackson PR, 88
Janowsky EC, 63
Johanson R, 69-70
 Johnston M, 110
Johnston SC, 91
Jones A, 56
Jones JIW, 54
Jones PW, 21, 22
 Jover JA, 20
 Julian J, 17
- Kalliomäki M, 106
Katz P, 93
 Katzelnick DJ, 48, 49
Kearon C, 72
 Keller MB, 103
 Kendrick D, 66
 Keneally JP, 62
Kent D, 34
 Kenyon SL, 69, 70
 Kerlikowske K, 51
 Kerr CR, 112
Khan KS, 36
 Khan MA, 67
 Khan MH, 67
 Kheim HB, 107
 Kidd C, 65
- King J, 10
 King M, 28
Kirkman R, 64
 Kleijnen J, 98
 Kline JA, 72
 Knight JF, 62
 Knight M, 10
 Ko WF, 51
 Koelemay MJ, 109
Koretz RL, 4
 Krabbe PF, 31
Kreder HJ, 104
 Kuck KH, 14
 Kuschner WG, 73
- Lacasse Y, 73
Laine L, 51, 54, 92
 Lalvani A, 75
 Landefeld CS, 18
Langhorne P, 12
 Lankisch PG, 77
 Lansang MA, 82
 Laterre P-F, 81
 Lau J, 115
Laupacis A, A-17 (No. 2), 60
Lawrence VA, 62
 Lee A, 62
 Legemate DA, 109
 Lehnick D, 77
Leibovici L, 55
 Lemelin J, 116
 Lijmer JG, 109
 Lilienthal J, 14
 Lin FYC, 107
 Lindström J, 101
 Lipsky PE, 3
 Little SH, 58
 Liu AH, 34
 Lloyd M, 28
 Lofaso F, 37
 Löfdahl CG, 23
 Løge AD, 12
 Lopez Bernaldo de Quiros JC, 26
 Luchetti R, 25
 Lundell L, 93
 Lung Health Study Research Group, 21
- Machin SJ, 96
 Maclean CC, 73
 MacLennan A, 105
Mahon JL, 41
 Majid M, 55
 Malarone International Study Team, 27
Malenka DJ, 32
 Mallorca FG, 71
 Malmivaara A, 104
Mangrulkar RS, A-14 (No. 3)
 Manning WG, 49
 Mansour D, 64
 Mantaring JB, 82
 Martin RW, 2
 Massel D, 58
- Mazur LJ, 24**
 Mazurski EJ, 44
 McCarty D, 17
McIntosh A, A-14 (No. 2)
McIvor A, 26
 McNamara RL, 13, 59
Meisner JS, 58
 Mewell JN, 67
 Miao YM, 33
 Michelson EA, 72
 Miettinen P, 93
 Miller MR, 13, 59
 Milne S, 99
 Miro JM, 26
Mitchell G, A-15 (No. 2)
 Mogenson CE, 9
 Monto AS, 42, 43
Montori VM, 45, 101
 Morado IC, 20
 Mørch B, 12
Mulrow CD, 6-7, 11
 Myrvold HE, 93
- Neer RM, 95
 Neldam S, 9
Neugut AI, 33
 Newell JN, 67
 Nguyen HC, 15
Nicolle LE, 83
Nidiry JJ, 77
Nikodem C, 65
 Norris SL, 45
 Nye FJ, 46
- O'Brien B, 19**
 O'Brien BJ, 113
O'Byrne PM, 23
O'Connor A, A-11-12 (No. 1)
 O'Meara SM, 55
 Olin J, 50
 ORACLE Collaborative Group, 69, 70
 Ormiston T, 87
 Oseltamivir Post Exposure Prophylaxis
 Investigator Group, 43
 Otto G, 76
 Owens DK, 73
- Packer M, 85
 Pahor M, 7
 Papi C, 25
 Paradise JL, 108
 Parish DC, 117
 Parkes J, 19
Pavia AT, 107
 Pearson SD, 48
 Peat JK, 44
Pelouze G-A, 110
 Peña JM, 26
Peters J, 22, 87
 Petrie KJ, 97
Peveler R, 48-49
Philbrick JT, 16

- Philp I, 111
 PLAF Investigators, 14
 Pomeroy IM, 111
 Postma DS, 23
 Powell P, 46
 Poynard T, 53
 Prins JB, 47
 Prins MH, 16
Pronovost PJ, 1
 Psaty BM, 7
Puntis J, 35

 Quinn L, 18

 Ramsay LE, 88
Rao G, 84-85
 Recombinant Human Activated Protein C
 Worldwide Evaluation in Severe Sepsis
 (PROWESS) Study Group, 81
 Regimbeau C, 53
 Ringash J, 63
 Rissanen P, 36
 Roalfe A, 114
Robbins JA, 100
 Roberts W, 65
 Robertson MC, 100
 Rodgers A, 1
 Roelandt JR, 32
 Rovers MM, 31
 Rubins HB, 86
Russell JA, 81
 Rydzak CE, 73

 Salminen S, 106
 Salpeter E, 87
 Salpeter S, 87
 Sanmuganathan PS, 88
Santos-Neto L, 68
 Scanlan J, 74
 Schaffner A, 83
Schmitt BP, 17
 Schneider L, 50

 Schoenfeld, 54
 Schug S, 1
 Scurr JH, 96
 Segal JB, 13, 59
 Seifert B, 83
 Seitsalo S, 104
Sellors J, A-19 (No. 1)
 Serruys PW, 89
 Shah S, 44
Shapiro CM, 97
Sharpe M, 46-47
Shehab T, 52
 Sheldon TA, 55
 Shelton RC, 103
 Shepperd S, 19
Sherman S, 111
 Shiffman ML, 52
 SHOCK Investigators, 57
Shoor S, 20
Silver A, 13
Simel DL, A-14 (No. 3)
 Simon GE, 48, 49
Simor AE, 75
 Sleeper LA, 57
 Sorrentino JV, 30
 Sousa JE, 89
 Sperber SJ, 30
 St. Clair EW, 3
 Stefanski K, 35
 Stevens JC, 94
 Stoker J, 109
 Straatman H, 31
 Streiner DL, 102
 Suen BY, 92

Talley NJ, 114
 Tarnow-Mordi W, 69, 70
 Tasche MJ, 24
 Tattersfield AE, 23
 Taylor DJ, 69, 70
 Taylor FC, 60
 Teperi J, 36

Thomas E, 115
 Tikkanen I, 9
Tilburt JC, A-14 (No. 3)
 Torkki M, 104
 Tuomilehto J, 101
 Turnbull D, 105
 Turner RB, 30

Uhari M, 71
 Uijen JH, 24
 Unger F, 89

 VA-HIT Study Group, 86
 van der Heide DM, 3
 van der Heijden JF, 16
 van der Wouden JC, 24
 van Walraven G, 117
 Vander Leek TK, 35
 Vargas F, 56
 Venkat Narayan KM, 45
 Vincent J-L, 81

 Walker N, 1
 Walley JD, 67
 Wallis EJ, 88
 Ward E, 28
 Welch V, 99
 Welliver R, 43
 Wheeler-Sherman, 71
 White HD, 57
Williams HC, 106
 Wilson S, 114

 Young LT, 102

 Zanamavir Family Study Group, 42
 Zanchetta JR, 95
 Zerah-Lancner F, 37
 Zuniga-Macaraig ZO, 71

A

- Abdominal pain
 - Review: Smooth-muscle relaxants improve symptoms and reduce pain in the irritable bowel syndrome, 53
- Accidental falls
 - A home-based, nurse-delivered exercise program reduced falls and serious injuries in persons ≥ 80 years of age, 100
- Acetamides
 - Oseltamivir reduced new cases of influenza in household contacts of persons with clinical or confirmed influenza, 43
- Acetaminophen
 - Bucindolol reduced mortality and hospitalization related to cardiovascular causes in advanced chronic heart failure, 84
 - Pseudoephedrine and acetaminophen relieved sinus symptoms in the course of the common cold, 30
- Adrenergic beta-antagonists
 - Carvedilol reduced mortality and hospitalization in severe chronic heart failure, 85
- AIDS-related opportunistic infections
 - Discontinuation of prophylaxis for *Pneumocystis carinii* pneumonia was safe after highly active antiretroviral therapy, 26
- Aircraft
 - Review: Oral melatonin reduces jet lag in air travelers, 97
- Albuminuria
 - Review: ACE inhibitors are beneficial in type 1 diabetes mellitus and microalbuminuria regardless of baseline risk factor status, 41
- Algorithms
 - The Mini-Cog had high sensitivity and specificity for diagnosing dementia in community-dwelling older adults, 74
- Alzheimer disease
 - Review: Galantamine improves most outcomes in suspected Alzheimer disease, 50
- Amiodarone
 - The implantable cardioverter defibrillator was not as cost-effective as amiodarone for prolonging survival, 113
 - Rate and rhythm control showed similar symptom improvement in atrial fibrillation, 14
- Amoxicillin-potassium clavulanate combination
 - Antibiotics did not prevent adverse outcomes after preterm, prelabor rupture of fetal membranes, 69
 - Antibiotics did not prevent adverse outcomes in women in preterm labor with intact fetal membranes, 70
- Amphotericin B
 - Continuous infusion of amphotericin B deoxycholate reduced overall mortality, side effects, and nephrotoxicity more than did rapid infusion, 83
- Analgesia
 - Incidence of long-term backache was similar between women who received epidural analgesia during labor and those who did not, 65
 - Review: Epidural or spinal anesthesia reduces postoperative mortality and morbidity, 1
 - Warm-water bathing did not reduce use of pharmacologic analgesia during the first stage of labor, 105
- Angioplasty
 - Percutaneous transluminal coronary angioplasty and endarterectomy were both effective for carotid stenosis, 91
 - Review: Glycoprotein IIb/IIIa inhibitors reduce MI and urgent revascularization but increase bleeding in PCI, 90
 - Stents had similar clinical outcomes but more repeat revascularization than did bypass surgery in multivessel disease, 89
- Angiotensin-converting enzyme inhibitors
 - Review: ACE inhibitors are beneficial in type 1 diabetes mellitus and microalbuminuria regardless of baseline risk factor status, 41
- Antagonists, vitamin K
 - Review: Low-molecular-weight heparin may be as effective and safe as vitamin-K antagonists in symptomatic venous thromboembolism, 16
- Anti-arrhythmic agents
 - The implantable cardioverter defibrillator was not as cost-effective as amiodarone for prolonging survival, 113
 - Review: Many drugs are effective for conversion of AF and maintenance of sinus rhythm, 13
- Antibiotics
 - Antibiotics did not prevent adverse outcomes after preterm, prelabor rupture of fetal membranes, 69
 - Antibiotics did not prevent adverse outcomes in women in preterm labor with intact fetal membranes, 70
- Antibodies, monoclonal
 - Infliximab plus methotrexate was more effective than methotrexate alone in rheumatoid arthritis, 3
- Anticoagulants
 - Multicomponent comprehensive care improved control and may have decreased major bleeding in older patients on warfarin, 18
 - Review: Anticoagulants may be better than antiplatelet agents for nonfatal stroke but not other vascular or fatal events in nonrheumatic AF, 60
 - Risk for a major bleed was higher for women than men on warfarin in patients with atrial fibrillation, 112
 - Vitamin K lowered the international normalized ratio into the therapeutic range in patients receiving warfarin, 17
- Antidepressants
 - Review: Antidepressants plus benzodiazepines lead to fewer dropouts and less depression severity in major depression, 102
- Antihypertensive agents
 - Review: ACE inhibitors, calcium antagonists, and more intensive blood pressure-lowering strategies reduce cardiovascular events, 6
 - Review: Calcium antagonists lead to a higher risk for MI, congestive heart failure, and major cardiovascular events, 7
- Anti-infective agents
 - Omeprazole was better than *H. pylori* eradication for preventing recurrent GI bleeding in patients receiving naproxen, 92
 - Permethrin (PER) plus trimethoprim/sulfamethoxazole (TMP/SMX) was better than PER but not TMP/SMX for treating head lice, 71
- Anti-inflammatory agents
 - Omeprazole was better than *H. pylori* eradication for preventing recurrent GI bleeding in patients receiving naproxen, 92
 - Recombinant human activated protein C reduced all-cause mortality in patients with severe sepsis, 81
- Antilipemic agents
 - Gemfibrozil reduced the risk for stroke in men with coronary heart disease and low levels of HDL cholesterol, 86
- Antimalarials
 - Malaria prophylaxis with atovaquone-proguanil caused fewer gastrointestinal adverse events than did chloroquine-proguanil, 27

- Mefloquine prevents malaria but has adverse effects that limit its acceptability in adults not immune to malarial infection, 68
- Antirheumatic agents**
- Etanercept was more effective and safer than methotrexate in reducing disease progression in early rheumatoid arthritis, 2
 - Infliximab plus methotrexate was more effective than methotrexate alone in rheumatoid arthritis, 3
- Antilulcer agents**
- Antireflux surgery may be more effective than low-dose, but not high-dose, omeprazole for gastroesophageal reflux disease, 93
- Antiviral agents**
- High-dose peginterferon α -2a sustained virologic and biochemical response in chronic hepatitis C infection with cirrhosis, 52
 - Oseltamivir reduced new cases of influenza in household contacts of persons with clinical or confirmed influenza, 43
 - Postexposure prophylaxis with inhaled zanamivir was efficacious for household contacts of persons with influenza, 42
- Arterial occlusive diseases**
- Review: Magnetic resonance angiography is highly accurate for diagnosing lower-extremity arterial disease in patients with claudication or critical limb ischemia, 109
- Arthritis rheumatoid**
- Etanercept was more effective and safer than methotrexate in reducing disease progression in early rheumatoid arthritis, 2
 - Infliximab plus methotrexate was more effective than methotrexate alone in rheumatoid arthritis, 3
- Aspirin**
- Aspirin but not vitamin E prevented cardiovascular events in patients at risk, 8
 - Clopidogrel reduced recurrent ischemic events in patients with previous cardiac surgery more than did aspirin, 5
 - Review: Anticoagulants may be better than antiplatelet agents for nonfatal stroke but not other vascular or fatal events in nonrheumatic AF, 60
 - Review: Antiplatelet drugs reduce preeclampsia, preterm birth, and stillbirth or neonatal death, 10
 - Review: Aspirin reduces the incidence of coronary artery disease in persons at risk, 88
- Asthma**
- Formoterol was more effective than terbutaline when taken as needed for moderate-to-severe asthma, 23
 - A peer-led asthma education program in adolescents was more effective than no program for improving quality of life, 44
 - Review: Inhaled beclomethasone reduces airflow limitations, symptoms, and the need for rescue bronchodilators in chronic asthma, 22
 - Review: Single-dose and longer-duration cardioselective β -blockers do not increase respiratory symptoms in reversible airway disease, 87
 - Review: Small treatment effect of disodium cromoglycate in childhood asthma is tempered by methodologic flaws and publication bias, 24
- Atherosclerosis**
- Aspirin but not vitamin E prevented cardiovascular events in patients at risk, 8
- Atrial fibrillation**
- Rate and rhythm control showed similar symptom improvement in atrial fibrillation, 14
 - Review: Anticoagulants may be better than antiplatelet agents for nonfatal stroke but not other vascular or fatal events in non-rheumatic AF, 60
 - Review: Many drugs are effective for conversion of AF and maintenance of sinus rhythm, 13
 - Risk for a major bleed was higher for women than men on warfarin in patients with atrial fibrillation, 112

B

- Back pain**
- Incidence of long-term backache was similar between women who received epidural analgesia during labor and those who did not, 65
- Bacterial infections**
- A prediction guide for patients with diarrhea detected non-*C. difficile* diarrhea and reduced culturing of stools, 75
- Bacterial polysaccharides**
- Salmonella typhi* Vi conjugate vaccine reduced the incidence of typhoid fever in 2- to 5-year-old children, 107
- Bandages**
- Wearing elastic compression stockings during long-haul flights prevented the development of deep venous thrombosis, 96
- Baths**
- Warm-water bathing did not reduce use of pharmacologic analgesia during the first stage of labor, 105
- Bayes theorem**
- Which methods for bedside Bayes? (Editorial), A-11(No.3)
- Beclomethasone**
- Review: Inhaled beclomethasone reduces airflow limitations, symptoms, and the need for rescue bronchodilators in chronic asthma, 22
- Benzodiazepines**
- Review: Antidepressants plus benzodiazepines lead to fewer dropouts and less depression severity in major depression, 102
- Beta-blocker, adrenergic**
- Review: Single-dose and longer-duration cardioselective β -blockers do not increase respiratory symptoms in reversible airway disease, 87
- Body mass index**
- Low Braden scale scores predicted the development of pressure ulcers in neurologic intensive and intermediate care units, 76
- Bone density**
- Parathyroid hormone decreased fracture rates and increased bone mineral density in postmenopausal women, 95
- Breast neoplasms**
- Review: Insufficient evidence exists for screening mammography in women 40 to 49 years of age, 63
- Budesonide**
- Review: Budesonide is better than mesalazine but not conventional corticosteroids for inducing remission in active Crohn disease, 25
- Bupivacaine**
- Incidence of long-term backache was similar between women who received epidural analgesia during labor and those who did not, 65

C

- Calcium channel blockers**
- Review: Calcium antagonists lead to a higher risk for MI, congestive heart failure, and major cardiovascular events, 7
- Cardiopulmonary resuscitation**
- A clinical decision aid was accurate for predicting survival to hospital discharge after in-hospital cardiac resuscitation, 117
- Cardiovascular diseases**
- Review: Aspirin reduces the incidence of coronary artery disease in persons at risk, 88

- Carotid stenosis
 Percutaneous transluminal coronary angioplasty and endarterectomy were both effective for carotid stenosis, 91
- CD4 lymphocyte count
 Discontinuation of prophylaxis for *Pneumocystis carinii* pneumonia was safe after highly active antiretroviral therapy, 26
- Cerebrovascular disorders
 Extended stroke-unit service improved independence at 26 weeks after acute stroke, 12
 Gemfibrozil reduced the risk for stroke in men with coronary heart disease and low levels of HDL cholesterol, 86
 Percutaneous transluminal coronary angioplasty and endarterectomy were both effective for carotid stenosis, 91
 Review: ACE inhibitors, calcium antagonists, and more intensive blood pressure-lowering strategies reduce cardiovascular events, 6
 Review: Warfarin prevents stroke in nonrheumatic atrial fibrillation but has a higher risk for hemorrhage than other agents, 59
 Stents had similar clinical outcomes but more repeat revascularization than did bypass surgery in multivessel disease, 89
- Child development
 No difference exists in developmental outcomes at 3 years of age between early or delayed tympanostomy for otitis media, 108
- Clopidogrel
 Clopidogrel reduced recurrent ischemic events in patients with previous cardiac surgery more than did aspirin, 5
- Clostridium difficile*
 A prediction guide for patients with diarrhea detected non-*C. difficile* diarrhea and reduced culturing of stools, 75
- Cognitive therapy
 Psychological therapy reduced depression earlier (4 months) but at 1 year was not better than usual general-practitioner care, 28
- Colonic diseases, functional
 Review: Smooth-muscle relaxants improve symptoms and reduce pain in the irritable bowel syndrome, 53
- Colonoscopy
 CT pneumocolon had a low sensitivity for detecting colorectal neoplasms, 33
- Colorectal neoplasms
 CT pneumocolon had a low sensitivity for detecting colorectal neoplasms, 33
- Common cold
 Pseudoephedrine and acetaminophen relieved sinus symptoms in the course of the common cold, 30
- Contraception
 Review: The levonorgestrel-releasing intrauterine system prevents more pregnancies than intrauterine devices ≤ 250 mm² but not IUDs > 250 mm², 64
- Coronary artery bypass
 Stents had similar clinical outcomes but more repeat revascularization than did bypass surgery in multivessel disease, 89
- Coronary disease
 Review: ACE inhibitors, calcium antagonists, and more intensive blood pressure-lowering strategies reduce cardiovascular events, 6
 Review: Aspirin reduces the incidence of coronary artery disease in persons at risk, 88
 Review: Dobutamine stress myocardial perfusion imaging has high sensitivity for detecting coronary artery disease, 32
 Review: Functional testing for restenosis after PTCA has limited value, 34
 Review: Glycoprotein IIb/IIIa inhibitors reduce MI and urgent revascularization but increase bleeding in PCI, 90
 Stents had similar clinical outcomes but more repeat revascularization than did bypass surgery in multivessel disease, 89
- Cost-benefit analysis
Helicobacter pylori testing and endoscopy was less cost-effective than usual management for patients with dyspepsia, 114
 A levonorgestrel-releasing intrauterine system was more cost-effective than was hysterectomy for menorrhagia, 36
- Crohn disease
 Review: Budesonide is better than mesalazine but not conventional corticosteroids for inducing remission in active Crohn disease, 25
- Cromolyn sodium
 Review: Small treatment effect of disodium cromoglycate in childhood asthma is tempered by methodologic flaws and publication bias, 24
- ## D
- Decision making
 Using patient decision aids to promote evidence-based decision making (Editorial), A-11-12 (No. 1)
- Decision support techniques
 A clinical decision aid was accurate for predicting survival to hospital discharge after in-hospital cardiac resuscitation, 117
- Decubitus ulcer
 Low Braden scale scores predicted the development of pressure ulcers in neurologic intensive and intermediate care units, 76
 Review: Some but not all topical antimicrobial agents improve the rate of healing of chronic wounds, 55
- Defibrillators, implantable
 The implantable cardioverter defibrillator was not as cost-effective as amiodarone for prolonging survival, 113
- Dementia
 The Mini-Cog had high sensitivity and specificity for diagnosing dementia in community-dwelling older adults, 74
 Review: Pharmacologic and nonpharmacologic interventions improve outcomes in patients with dementia and for their caregivers, 94
- Deoxycholic acid
 Continuous infusion of amphotericin B deoxycholate reduced overall mortality, side effects, and nephrotoxicity more than did rapid infusion, 83
- Depression
 A depression-management program increased depression-free days and costs in depressed frequent users of general health care, 49
 A depression-management program reduced depression in frequent users of health care but did not reduce health care visits, 48
 Psychological therapy reduced depression earlier (4 months) but at 1 year was not better than usual general-practitioner care, 28
 Review: Antidepressants plus benzodiazepines lead to fewer dropouts and less depression severity in major depression, 102
 Short scales were as effective as long scales in screening for depression in older patients, 111
 St. John's wort was not better than placebo for reducing depression scores, 103
- Dermatitis, atopic
 Pre- and postnatal administration of *Lactobacillus* GG reduced the occurrence of atopic disease in offspring, 106

- Diabetes mellitus**
 Changes in diet and physical activity prevented type 2 diabetes mellitus in persons with impaired glucose tolerance, 101
 Combination therapy with candesartan and lisinopril was more effective than monotherapy in type 2 diabetes and hypertension, 9
 Edema reduction by mechanical compression improved the healing of foot infection in patients with diabetes mellitus, 15
 Review: ACE inhibitors are beneficial in type 1 diabetes mellitus and microalbuminuria regardless of baseline risk factor status, 41
 Review: Self-management training in type 2 diabetes mellitus is effective in the short term, 45
- Diabetic nephropathies**
 Review: ACE inhibitors are beneficial in type 1 diabetes mellitus and microalbuminuria regardless of baseline risk factor status, 41
- Diagnosis, differential**
 A 2-step procedure determined diagnosis in patients with unexplained weight loss, 77
- Diarrhea**
 A prediction guide for patients with diarrhea detected non-*C. difficile* diarrhea and reduced culturing of stools, 75
- Diet**
 Changes in diet and physical activity prevented type 2 diabetes mellitus in persons with impaired glucose tolerance, 101
 Reduced sodium intake lowered blood pressure and need for antihypertensive medication, 61
- Diltiazem**
 Rate and rhythm control showed similar symptom improvement in atrial fibrillation, 14
- Diphtheria-tetanus-acellular pertussis vaccines**
 Routine primary immunization using a longer needle resulted in fewer local reactions in infants, 29
- Disease transmission**
 Postexposure prophylaxis with inhaled zanamivir was efficacious for household contacts of persons with influenza, 42
- Dobutamine**
 Review: Dobutamine stress myocardial perfusion imaging has high sensitivity for detecting coronary artery disease, 32
- Drug administration schedule**
 Continuous infusion of amphotericin B deoxycholate reduced overall mortality, side effects, and nephrotoxicity more than did rapid infusion, 83
 Direct observation of tuberculosis treatment did not promote higher cure rates than did self-administered treatment, 67
- Drug monitoring**
 Multicomponent comprehensive care improved control and may have decreased major bleeding in older patients on warfarin, 18
- Dyspepsia**
Helicobacter pylori testing and endoscopy was less cost-effective than usual management for patients with dyspepsia, 114
 Review: *Helicobacter pylori* therapy does not reduce symptoms in nonulcer dyspepsia, 54
- E**
- Echocardiography**
 Review: Dobutamine stress myocardial perfusion imaging has high sensitivity for detecting coronary artery disease, 32
- Edema**
 Edema reduction by mechanical compression improved the healing of foot infection in patients with diabetes mellitus, 15
- Education, medical, graduate**
 Creating knowledge management skills in primary care residents: a description of a new pathway to evidence-based practice (Editorial), A-11-12 (No. 2)
- Electrocardiography**
 Review: Functional testing for restenosis after PTCA has limited value, 34
- Endarterectomy**
 Percutaneous transluminal coronary angioplasty and endarterectomy were both effective for carotid stenosis, 91
- Enzyme inhibitors**
 Antireflux surgery may be more effective than low-dose, but not high-dose, omeprazole for gastroesophageal reflux disease, 93
 Oseltamivir reduced new cases of influenza in household contacts of persons with clinical or confirmed influenza, 43
- Ephedrine**
 Pseudoephedrine and acetaminophen relieved sinus symptoms in the course of the common cold, 30
- Erythromycin**
 Antibiotics did not prevent adverse outcomes after preterm, prelabor rupture of fetal membranes, 69
 Antibiotics did not prevent adverse outcomes in women in preterm labor with intact fetal membranes, 70
- Esophageal and gastric varices**
 Review: Octreotide reduces rebleeding more than does vasopressin or terlipressin in patients with esophageal varices, 51
- Evidence-based medicine**
 Creating knowledge management skills in primary care residents: a description of a new pathway to evidence-based practice (Editorial), A-11-12 (No. 2)
Evidence-Based On Call Acute Medicine (Resource Corner), A-14 (No. 3)
Evidence-Based Physical Diagnosis (Resource Corner), A-14 (No. 3)
Evidence-Based Practice: A Primer for Health Care Professionals (Resource Corner), A-19 (No. 1)
An Evidence-Based Resource for Pain Relief (Resource Corner), A-15 (No. 2)
 Which methods for bedside Bayes? (Editorial), A-11 (No. 3)
- Exercise**
 Changes in diet and physical activity prevented type 2 diabetes mellitus in persons with impaired glucose tolerance, 101
 A home-based, nurse-delivered exercise program reduced falls and serious injuries in persons ≥ 80 years of age, 100
 Patient education to encourage graded exercise improved physical functioning in the chronic fatigue syndrome, 46
- F**
- Family practice**
 A tailored, multifaceted program in capitation-based family practices improved appropriate use of preventive-care maneuvers, 116
- Fatigue syndrome, chronic**
 Cognitive behavior therapy reduced fatigue severity and functional impairment in the chronic fatigue syndrome, 47
 Patient education to encourage graded exercise improved physical functioning in the chronic fatigue syndrome, 46
- Feces**
 A prediction guide for patients with diarrhea detected non-*C. difficile* diarrhea and reduced culturing of stools, 75

- Fetal membranes, premature rupture
 - Antibiotics did not prevent adverse outcomes after preterm, prelabor rupture of fetal membranes, 69
- Fibrin fibrinogen degradation products
 - A negative quantitative latex D-dimer assay helped to rule out venous thromboembolism, 110
 - A normal alveolar dead-space fraction plus D-dimer assay ruled out suspected pulmonary embolism in the emergency department, 72
- Fibrinolytic agents
 - Recombinant human activated protein C reduced all-cause mortality in patients with severe sepsis, 81
- Foot ulcer, diabetic
 - Review: Some but not all topical antimicrobial agents improve the rate of healing of chronic wounds, 55
- Formoterol
 - Formoterol was more effective than terbutaline when taken as needed for moderate-to-severe asthma, 23
- Fractures
 - Parathyroid hormone decreased fracture rates and increased bone mineral density in postmenopausal women, 95

G

- Galantamine
 - Review: Galantamine improves most outcomes in suspected Alzheimer disease, 50
- Gastroesophageal reflux
 - Antireflux surgery may be more effective than low-dose, but not high-dose, omeprazole for gastroesophageal reflux disease, 93
- Gastrointestinal disorders
 - Malaria prophylaxis with atovaquone-proguanil caused fewer gastrointestinal adverse events than did chloroquine-proguanil, 27
 - Omeprazole was better than *H. pylori* eradication for preventing recurrent GI bleeding in patients receiving naproxen, 92
 - Review: Octreotide reduces rebleeding more than does vasopressin or terlipressin in patients with esophageal varices, 51
 - A 2-step procedure determined diagnosis in patients with unexplained weight loss, 77
- Geriatric assessment
 - Short scales were as effective as long scales in screening for depression in older patients, 111
- Glucocorticoids
 - Inhaled triamcinolone did not slow the decline in pulmonary function in patients with COPD, 21
 - Review: Budesonide is better than mesalazine but not conventional corticosteroids for inducing remission in active Crohn disease, 25
- Glucose intolerance
 - Changes in diet and physical activity prevented type 2 diabetes mellitus in persons with impaired glucose tolerance, 101

H

- Haemophilus* vaccines
 - Routine primary immunization using a longer needle resulted in fewer local reactions in infants, 29
- Hallux valgus
 - Surgery was more effective than orthosis for hallux valgus, 104
- Health resources
 - A depression-management program increased depression-free days and costs in depressed frequent users of general health care, 49

- A depression-management program reduced depression in frequent users of health care but did not reduce health care visits, 48
- Heart failure
 - Bucindolol reduced mortality and hospitalization related to cardiovascular causes in advanced chronic heart failure, 84
 - Carvedilol reduced mortality and hospitalization in severe chronic heart failure, 85
 - A clinical decision aid was accurate for predicting survival to hospital discharge after in-hospital cardiac resuscitation, 117
- Helicobacter pylori*
 - Helicobacter pylori* testing and endoscopy was less cost-effective than usual management for patients with dyspepsia, 114
 - Omeprazole was better than *H. pylori* eradication for preventing recurrent GI bleeding in patients receiving naproxen, 92
 - Review: *Helicobacter pylori* therapy does not reduce symptoms in nonulcer dyspepsia, 54
- Hemorrhage
 - Multicomponent comprehensive care improved control and may have decreased major bleeding in older patients on warfarin, 18
 - Review: Warfarin prevents stroke in nonrheumatic atrial fibrillation but has a higher risk for hemorrhage than other agents, 59
- Heparin, low-molecular-weight
 - Review: Low-molecular-weight heparin may be as effective and safe as vitamin-K antagonists in symptomatic venous thromboembolism, 16
- Hepatitis C, chronic
 - High-dose peginterferon α -2a sustained virologic and biochemical response in chronic hepatitis C infection with cirrhosis, 52
- Hepatitis, alcoholic
 - Pentoxifylline improved short-term survival in severe alcoholic hepatitis, 4
- HIV infections
 - Discontinuation of prophylaxis for *Pneumocystis carinii* pneumonia was safe after highly active antiretroviral therapy, 26
- Home care services
 - A home-based, nurse-delivered exercise program reduced falls and serious injuries in persons ≥ 80 years of age, 100
 - Extended stroke-unit service improved independence at 26 weeks after acute stroke, 12
- Hospital units
 - Extended stroke-unit service improved independence at 26 weeks after acute stroke, 12
 - How to Read a Paper* (Resource Corner), A-14 (No. 2)
- Hypericum
 - St. John's wort was not better than placebo for reducing depression scores, 103
- Hypersensitivity
 - Routine primary immunization using a longer needle resulted in fewer local reactions in infants, 29
- Hypertension
 - Combination therapy with candesartan and lisinopril was more effective than monotherapy in type 2 diabetes and hypertension, 9
 - Reduced sodium intake lowered blood pressure and need for antihypertensive medication, 61
 - Review: Evidence is sparse and inconclusive for treating and monitoring chronic mild-to-moderate hypertension in pregnancy, 11
- Hysterectomy
 - A levonorgestrel-releasing intrauterine system was more cost-effective than was hysterectomy for menorrhagia, 36

I

- Immunization
Routine primary immunization using a longer needle resulted in fewer local reactions in infants, 29
- Immunocompromised host
Noninvasive intermittent ventilation reduced intubation and mortality in immunosuppressed patients with acute respiratory failure, 56
- Immunoglobulins
Review: Intravenous polyclonal immunoglobulins reduce mortality in patients with bacterial sepsis and septic shock, 82
Review: Glycoprotein IIb/IIIa inhibitors reduce MI and urgent revascularization but increase bleeding in PCI, 90
- Indoprofen
Review: Anticoagulants may be better than antiplatelet agents for nonfatal stroke but not other vascular or fatal events in nonrheumatic AF, 60
- Influenza
Oseltamivir reduced new cases of influenza in household contacts of persons with clinical or confirmed influenza, 43
Postexposure prophylaxis with inhaled zanamivir was efficacious for household contacts of persons with influenza, 42
- Insecticides
Permethrin (PER) plus trimethoprim/sulfamethoxazole (TMP/SMX) was better than PER but not TMP/SMX for treating head lice, 71
- Intensive care units
Low Braden scale scores predicted the development of pressure ulcers in neurologic intensive and intermediate care units, 76
- Interferon alfa-2a
High-dose peginterferon α -2a sustained virologic and biochemical response in chronic hepatitis C infection with cirrhosis, 52
- International normalized ratio
Vitamin K lowered the international normalized ratio into the therapeutic range in patients receiving warfarin, 17
- Intrauterine devices, medicated
Review: The levonorgestrel-releasing intrauterine system prevents more pregnancies than intrauterine devices ≤ 250 mm² but not IUDs > 250 mm², 64

L

- Labor
Antibiotics did not prevent adverse outcomes after preterm, prelabor rupture of fetal membranes, 69
Antibiotics did not prevent adverse outcomes in women in preterm labor with intact fetal membranes, 70
Warm-water bathing did not reduce use of pharmacologic analgesia during the first stage of labor, 105
- Lactobacillus
Pre- and postnatal administration of *Lactobacillus* GG reduced the occurrence of atopic disease in offspring, 106
- Latex fixation tests
A negative quantitative latex D-dimer assay helped to rule out venous thromboembolism, 110
- Leg ulcer
Review: Some but not all topical antimicrobial agents improve the rate of healing of chronic wounds, 55
- Levonorgestrel
A levonorgestrel-releasing intrauterine system was more cost-effective than was hysterectomy for menorrhagia, 36

Review: The levonorgestrel-releasing intrauterine system prevents more pregnancies than intrauterine devices ≤ 250 mm² but not IUDs > 250 mm², 64

- Lice infestations
Permethrin (PER) plus trimethoprim/sulfamethoxazole (TMP/SMX) was better than PER but not TMP/SMX for treating head lice, 71
- Lipoproteins, HDL cholesterol
Gemfibrozil reduced the risk for stroke in men with coronary heart disease and low levels of HDL cholesterol, 86
- Lisinopril
Combination therapy with candesartan and lisinopril was more effective than monotherapy in type 2 diabetes and hypertension, 9
- Liver cirrhosis
High-dose peginterferon α -2a sustained virologic and biochemical response in chronic hepatitis C infection with cirrhosis, 52
- Low-back pain
Radiography led to improved patient satisfaction but increased short-term pain in patients with low-back pain, 66
Review: Transcutaneous electrical nerve stimulation is not effective for chronic low-back pain, 99
- Lung disease, obstructive
Inhaled triamcinolone did not slow the decline in pulmonary function in patients with COPD, 21
Review: Single-dose and longer-duration cardioselective β -blockers do not increase respiratory symptoms in reversible airway disease, 87

M

- Magnetic resonance angiography
Review: Magnetic resonance angiography is highly accurate for diagnosing lower-extremity arterial disease in patients with claudication or critical limb ischemia, 109
- Malaria
Malaria prophylaxis with atovaquone-proguanil caused fewer gastrointestinal adverse events than did chloroquine-proguanil, 27
Mefloquine prevents malaria but has adverse effects that limit its acceptability in adults not immune to malarial infection, 68
- Mammography
Review: Insufficient evidence exists for screening mammography in women 40 to 49 years of age, 63
- Mass screening
Review: Insufficient evidence exists for screening mammography in women 40 to 49 years of age, 63
- Medical errors
Review: Some interventions are effective in reducing medical errors, 115
- Mefloquine
Mefloquine prevents malaria but has adverse effects that limit its acceptability in adults not immune to malarial infection, 68
- Melatonin
Review: Oral melatonin reduces jet lag in air travelers, 97
- Menorrhagia
A levonorgestrel-releasing intrauterine system was more cost-effective than was hysterectomy for menorrhagia, 36
- Methotrexate
Etanercept was more effective and safer than methotrexate in reducing disease progression in early rheumatoid arthritis, 2
Infliximab plus methotrexate was more effective than methotrexate alone in rheumatoid arthritis, 3

- Methotrexate plus prednisone was more effective than prednisone monotherapy in giant-cell arteritis, 20
- Middle ear ventilation
Ventilation tubes did not improve quality of life in persistent otitis media with effusion, 31
- Myocardial infarction
Early revascularization improved 1-year survival after myocardial infarction with cardiogenic shock, 57
Review: Calcium antagonists lead to a higher risk for MI, congestive heart failure, and major cardiovascular events, 7
- Myocardial revascularization
Early revascularization improved 1-year survival after myocardial infarction with cardiogenic shock, 57

N

- Needles
Routine primary immunization using a longer needle resulted in fewer local reactions in infants, 29
- Neoplasm, pulmonary
Review: Positron emission tomography with 18-fluorodeoxyglucose is an accurate test for identifying malignant pulmonary lesions, 73
- Neuraminidase
Oseltamivir reduced new cases of influenza in household contacts of persons with clinical or confirmed influenza, 43
- Neurology
Review: Pharmacologic and nonpharmacologic interventions improve outcomes in patients with dementia and for their caregivers, 94
- Neuropsychological tests
The Mini-Cog had high sensitivity and specificity for diagnosing dementia in community-dwelling older adults, 74
- Neutropenia
Continuous infusion of amphotericin B deoxycholate reduced overall mortality, side effects, and nephrotoxicity more than did rapid infusion, 83

O

- Octreotide
Review: Octreotide reduces rebleeding more than does vasopressin or terlipressin in patients with esophageal varices, 51
- Omeprazole
Antireflux surgery may be more effective than low-dose, but not high-dose, omeprazole for gastroesophageal reflux disease, 93
- Orthotic devices
Surgery was more effective than orthosis for hallux valgus, 104
- Osteoporosis
Parathyroid hormone decreased fracture rates and increased bone mineral density in postmenopausal women, 95
- Osteotomy
Surgery was more effective than orthosis for hallux valgus, 104
- Otitis media with effusion
No difference exists in developmental outcomes at 3 years of age between early or delayed tympanostomy for otitis media, 108
Ventilation tubes did not improve quality of life in persistent otitis media with effusion, 31
- Oxygen inhalation therapy
Noninvasive intermittent ventilation reduced intubation and mortality in immunosuppressed patients with acute respiratory failure, 56

P

- Pain
An Evidence-Based Resource for Pain Relief (Resource Corner), A-15 (No. 2)
Warm-water bathing did not reduce use of pharmacologic analgesia during the first stage of labor, 105
- Parasympatholytics
Review: Smooth-muscle relaxants improve symptoms and reduce pain in the irritable bowel syndrome, 53
- Patient care planning
Multicomponent comprehensive care improved control and may have decreased major bleeding in older patients on warfarin, 18
- Patient discharge
Extended stroke-unit service improved independence at 26 weeks after acute stroke, 12
Review: Evidence of the effectiveness of discharge planning is equivocal, 19
- Patient education
Multicomponent comprehensive care improved control and may have decreased major bleeding in older patients on warfarin, 18
Patient education to encourage graded exercise improved physical functioning in the chronic fatigue syndrome, 46
A peer-led asthma education program in adolescents was more effective than no program for improving quality of life, 44
Review: Self-management training in type 2 diabetes mellitus is effective in the short term, 45
Using patient decision aids to promote evidence-based decision making (Editorial), A-11-12 (No. 1)
- Peanuts, hypersensitivity
Subsequent reactions were common and often more serious than the initial reactions of children with peanut allergy, 35
- Pediculus
Permethrin (PER) plus trimethoprim/sulfamethoxazole (TMP/SMX) was better than PER but not TMP/SMX for treating head lice, 71
- Peer group
A peer-led asthma education program in adolescents was more effective than no program for improving quality of life, 44
- Pentoxifylline
Pentoxifylline improved short-term survival in severe acute alcoholic hepatitis, 4
- Peptides
Review: Glycoprotein IIb/IIIa inhibitors reduce MI and urgent revascularization but increase bleeding in PCI, 90
- Peripheral vascular diseases
Review: Magnetic resonance angiography is highly accurate for diagnosing lower-extremity arterial disease in patients with claudication or critical limb ischemia, 109
- Physical examination
Evidence-Based Physical Diagnosis (Resource Corner), A-14 (No. 3)
- Physician's practice patterns
A tailored, multifaceted program in capitation-based family practices improved appropriate use of preventive-care maneuvers, 116
- Physician-patient relations
Review: Cognitive care and combined cognitive and emotional care interventions may influence patients' health outcomes, 98
- Platelet aggregation inhibitors
Clopidogrel reduced recurrent ischemic events in patients with previous cardiac surgery more than did aspirin, 5

- Review: Anticoagulants may be better than antiplatelet agents for nonfatal stroke but not other vascular or fatal events in nonrheumatic AF, 60
- Review: Antiplatelet drugs reduce preeclampsia, preterm birth, and stillbirth or neonatal death, 10
- Review: Glycoprotein IIb/IIIa inhibitors reduce MI and urgent revascularization but increase bleeding in PCI, 90
- Pneumocystis carinii*
- Discontinuation of prophylaxis for *Pneumocystis carinii* pneumonia was safe after highly active antiretroviral therapy, 26
- Pneumonia
- Discontinuation of prophylaxis for *Pneumocystis carinii* pneumonia was safe after highly active antiretroviral therapy, 26
- Pneumoradiography
- CT pneumocolon had a low sensitivity for detecting colorectal neoplasms, 33
- Point-of-care systems
- Helicobacter pylori* testing and endoscopy was less cost-effective than usual management for patients with dyspepsia, 114
- Polysomnography
- A predictive model using pulmonary-function parameters identified snorers at low risk for the sleep apnea syndrome, 37
- Positive-pressure respiration
- Noninvasive intermittent ventilation reduced intubation and mortality in immunosuppressed patients with acute respiratory failure, 56
- Postoperative complications
- Review: Epidural or spinal anesthesia reduces postoperative mortality and morbidity, 1
- Prednisone
- Methotrexate plus prednisone was more effective than prednisone monotherapy in giant-cell arteritis, 20
- Preeclampsia
- Review: Antiplatelet drugs reduce preeclampsia, preterm birth, and stillbirth or neonatal death, 10
- Review: Evidence is sparse and inconclusive for treating and monitoring chronic mild-to-moderate hypertension in pregnancy, 11
- Pregnancy
- Review: Evidence is sparse and inconclusive for treating and monitoring chronic mild-to-moderate hypertension in pregnancy, 11
- Preventive health services
- A tailored, multifaceted program in capitation-based family practices improved appropriate use of preventive-care maneuvers, 116
- Primary health care
- Creating knowledge management skills in primary care residents: a description of a new pathway to evidence-based practice (Editorial), A-11-12 (No. 2)
- Evidence-Based Practice: A Primer for Health Care Professionals (Resource Corner), A-19 (No. 1)
- A tailored, multifaceted program in capitation-based family practices improved appropriate use of preventive-care maneuvers, 116
- Probiotics
- Pre- and postnatal administration of *Lactobacillus* GG reduced the occurrence of atopic disease in offspring, 106
- Protein C
- Recombinant human activated protein C reduced all-cause mortality in patients with severe sepsis, 81
- Psychotherapy
- Cognitive behavior therapy reduced fatigue severity and functional impairment in the chronic fatigue syndrome, 47
- Pulmonary embolism
- A normal alveolar dead-space fraction plus D-dimer assay ruled out suspected pulmonary embolism in the emergency department, 72
- Pyrethrins
- Permethrin (PER) plus trimethoprim/sulfamethoxazole (TMP/SMX) was better than PER but not TMP/SMX for treating head lice, 71
- ## Q
- Quality of life
- Ventilation tubes did not improve quality of life in persistent otitis media with effusion, 31
- ## R
- Radiography
- Radiography led to improved patient satisfaction but increased short-term pain in patients with low-back pain, 66
- Radionuclide imaging
- Review: Functional testing for restenosis after PTCA has limited value, 34
- Reading
- How to Read a Paper* (Resource Corner), A-14 (No. 2)
- Receptors, tumor necrosis factor
- Etanercept was more effective and safer than methotrexate in reducing disease progression in early rheumatoid arthritis, 2
- Recombinant human activated protein C reduced all-cause mortality in patients with severe sepsis, 81
- Recombinant proteins
- Recombinant human activated protein C reduced all-cause mortality in patients with severe sepsis, 81
- Respiratory function tests
- A normal alveolar dead-space fraction plus D-dimer assay ruled out suspected pulmonary emboli in the emergency department, 72
- A predictive model using pulmonary-function parameters identified snorers at low risk for the sleep apnea syndrome, 37
- Respiratory insufficiency
- Noninvasive intermittent ventilation reduced intubation and mortality in immunosuppressed patients with acute respiratory failure, 56
- Resuscitation orders
- A clinical decision aid was accurate for predicting survival to hospital discharge after in-hospital cardiac resuscitation, 117
- Review: Anticoagulants may be better than antiplatelet agents for nonfatal stroke but not other vascular or fatal events in nonrheumatic AF (Letter), A-17 (No. 2)
- Review: Warfarin prevents stroke in nonrheumatic atrial fibrillation but has a higher risk for hemorrhage than other agents, 59
- ## S
- Salmonella typhi*
- Salmonella typhi* Vi conjugate vaccine reduced the incidence of typhoid fever in 2 to 5-year-old children, 107
- Scalp dermatoses
- Permethrin (PER) plus trimethoprim/sulfamethoxazole (TMP/SMX) was better than PER but not TMP/SMX for treating head lice, 71

Self care

Review: Self-management training in type 2 diabetes mellitus is effective in the short term, 45

Sepsis

Review: Intravenous polyclonal immunoglobulins reduce mortality in patients with bacterial sepsis and septic shock, 82

Sex factors

Risk for a major bleed was higher for women than men on warfarin in patients with atrial fibrillation, 112

Shock

Early revascularization improved 1-year survival after myocardial infarction with cardiogenic shock, 57

Review: Intravenous polyclonal immunoglobulins reduce mortality in patients with bacterial sepsis and septic shock, 82

Sialic acids

Postexposure prophylaxis with inhaled zanamivir was efficacious for household contacts of persons with influenza, 42

Sleep apnea syndromes

A predictive model using pulmonary-function parameters identified snorers at low risk for the sleep apnea syndrome, 37

Sleep disorders

Review: Oral melatonin reduces jet lag in air travelers, 97

Stents

Clopidogrel reduced recurrent ischemic events in patients with previous cardiac surgery more than did aspirin, 5

Review: Functional testing for restenosis after PTCA has limited value, 34

T

Tachycardia, ventricular

The implantable cardioverter defibrillator was not as cost-effective as amiodarone for prolonging survival, 113

Temporal arteritis

Methotrexate plus prednisone was more effective than prednisone monotherapy in giant-cell arteritis, 20

Terbutaline

Formoterol was more effective than terbutaline when taken as needed for moderate-to-severe asthma, 23

Thromboembolism

Review: Low-molecular-weight heparin may be as effective and safe as vitamin-K antagonists in symptomatic venous thromboembolism, 16

Tomography

CT pneumocolon had a low sensitivity for detecting colorectal neoplasms, 33

Review: Positron emission tomography with 18-fluorodeoxyglucose is an accurate test for identifying malignant pulmonary lesions, 73

Transcutaneous electric nerve stimulation

Review: Transcutaneous electrical nerve stimulation is not effective for chronic low-back pain, 99

Travel

Mefloquine prevents malaria but has adverse effects that limit its acceptability in adults not immune to malarial infection, 68

Review: Oral melatonin reduces jet lag in air travelers, 97

Wearing elastic compression stockings during long-haul flights prevented the development of deep venous thrombosis, 96

Triamcinolone

Inhaled triamcinolone did not slow the decline in pulmonary function in patients with COPD, 21

Trimethoprim-sulfamethoxazole combination

Permethrin (PER) plus trimethoprim/sulfamethoxazole (TMP/SMX) was better than PER but not TMP/SMX for treating head lice, 71

Tuberculosis

Direct observation of tuberculosis treatment did not promote higher cure rates than did self-administered treatment, 67

Tumor necrosis factor

Infliximab plus methotrexate was more effective than methotrexate alone in rheumatoid arthritis, 3

Tympanoplasty

No difference exists in developmental outcomes at 3 years of age between early or delayed tympanostomy for otitis media, 108

Typhoid

Salmonella typhi Vi conjugate vaccine reduced the incidence of typhoid fever in 2- to 5-year-old children, 107

V

Venous thrombosis

Multicomponent comprehensive care improved control and may have decreased major bleeding in older patients on warfarin, 18

A negative quantitative latex D-dimer assay helped to rule out venous thromboembolism, 110

Wearing elastic compression stockings during long-haul flights prevented the development of deep venous thrombosis, 96

Ventricular fibrillation

The implantable cardioverter defibrillator was not as cost-effective as amiodarone for prolonging survival, 113

Vitamins

Aspirin but not vitamin E prevented cardiovascular events in patients at risk, 8

Vitamin K lowered the international normalized ratio into the therapeutic range in patients receiving warfarin, 17

W

Warfarin

Multicomponent comprehensive care improved control and may have decreased major bleeding in older patients on warfarin, 18

Review: Calcium antagonists lead to a higher risk for MI, congestive heart failure, and major cardiovascular events, 7

Review: Warfarin prevents stroke in nonrheumatic atrial fibrillation but has a higher risk for hemorrhage than other agents, 59

Vitamin K lowered the international normalized ratio into the therapeutic range in patients receiving warfarin, 17

Weight loss

A 2-step procedure determined diagnosis in patients with unexplained weight loss, 77

Wound infection

Edema reduction by mechanical compression improved the healing of foot infection in patients with diabetes mellitus, 15

